

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 : A61K 45/00, 9/20, A61P 3/06	A1	(11) International Publication Number: WO 00/62810 (43) International Publication Date: 26 October 2000 (26.10.00)
(21) International Application Number: PCT/SE00/00755 (22) International Filing Date: 19 April 2000 (19.04.00) (30) Priority Data: 9901387-2 19 April 1999 (19.04.99) SE (71) Applicant (for all designated States except US): AS-TRAZENECA AB [SE/SE]; S-151 85 Södertälje (SE). (72) Inventors; and (75) Inventors/Applicants (for US only): ABRAHAMSSON, Bertil [SE/SE]; AstraZeneca R & D Mölndal, S-431 83 Mölndal (SE). LINDQVIST, Ann-Margret [SE/SE]; AstraZeneca R & D Mölndal, S-431 83 Mölndal (SE). UNGELL, Anna-Lena [SE/SE]; AstraZeneca R & D Mölndal, S-431 83 Mölndal (SE). (74) Agent: ASTRAZENECA AB; Global Intellectual Property, Patents, S-151 85 Södertälje (SE).		(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: AN ORAL FORMULATION FOR ILEUM ADMINISTERING COMPRISING AN INHIBITOR COMPOUND OF THE ILEAL BILE ACID TRANSPORT (57) Abstract An oral pharmaceutical formulation comprising an inhibitor compound of the ileal bile acid transport (IBAT inhibitor compound) and a therapeutically acceptable carrier characterised in that the formulation is designed to deliver the IBAT inhibitor compound in the ileum. The IBAT inhibitor compound can also be administered in combination with a bile acid binder to alleviate possible side effects of therapy with IBAT inhibitor compound, such as for instance diarrhoea. The bile acid binder may be formulated for colon release.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

AN ORAL FORMULATION FOR ILEUM ADMINISTERING COMPRISING AN INHIBITOR COMPOUND OF THE ILEAL BILE ACID TRANSPORT**FIELD OF THE INVENTION**

5 The present invention relates to an oral pharmaceutical dosage form comprising a substance with inhibiting effect on the ileal bile acid transport system (IBAT). More specifically, the dosage form is suitable in the treatment of hypercholesterolaemia. The invention also relates to manufacturing processes and the use of the dosage form in the treatment of hypercholesterolaemia. A further aspect of the invention is the use of a
10 substance with inhibiting effect on IBAT in combination with a bile acid binder by simultaneously, separately or sequentially administration of the two substances, and the use of these substances in the manufacture of such a pharmaceutical dosage form.

BACKGROUND OF THE INVENTION AND PRIOR ART

15 It is well known that hyperlipidemic conditions associated with elevated concentrations of total cholesterol and low-density lipoprotein cholesterol are major risk factors for coronary heart disease and particularly atherosclerosis. Interfering with the circulation of bile acids within the lumen of the intestinal tracts is found to reduce the level of cholesterol. Previous
20 established therapies to reduce the concentration of cholesterol involve for instance treatment with HMG-CoA reductase inhibitors, preferably statins such as simvastatin and fluvastatin, or treatment with bile acid binders, such as resins. Frequently used bile acid binders are for instance cholestyramine and cholestipol. One recently proposed therapy involves the treatment with substances with inhibiting effect on the ileal bile acid transport system
25 (IBAT).

Re-absorption of bile acid from the gastro-intestinal tract is a normal physiological process, which mainly takes place in the ileum by an active transport mechanism called ileal bile acid transport (IBAT). Inhibitors of IBAT can be used in the treatment of
30 hypercholesterolaemia. See for instance "Interaction of bile acids and cholesterol with

nonsystemic agents having hypocholesterolemic properties", *Biochemica et Biophysica Acta*, 1210 (1994) 255- 287. Thus, suitable compounds having such inhibitory IBAT activity are also useful in the treatment of hyperlipidaemic conditions.

- 5 Several chemical compounds possessing such IBAT activity have recently been described, see for instance hypolipidaemic benzothiazepine compounds described in International Patent Application, Publication No. WO 93/16055 and WO 96/16051; condensed 1,4-thiazepines described in International Patent Application, Publication No. WO 94/18183; different heterocyclic compounds described in International Patent Application,
10 Publication No. WO 94/18184; and 1,4-benzothiazepine-1,1-dioxides described in International Patent Application, Publication No. WO 96/05188, all of which are hereby incorporated by reference.

- Further, especially suitable compounds for the present invention are for instance
15 benzothiazepines with IBAT activity described in International Patent Application, Publication No. WO 96/08484; bile acid resorption inhibitors described in International Patent Application, Publication No. WO 97/33882, WO 98/07449 and WO 98/03818, and in European Patent Application, Publication No. EP-A-0864582, EP-A-0489423, EP-A-0549967, EP-A-0573848, EP-A-0624593, EP-A-0624594, EP-A-0624595, and EP-A-
20 0624596, all of which are hereby incorporated by reference. Further compounds of interest can be found in International Patent Application, Publication No. WO 99/32478, WO 99/64409 and WO 00/01687, all of which are hereby incorporated by reference.

- It is proposed that these types of compounds can be administered by any conventional
25 means available for use in conjunction with pharmaceuticals. For instance, the dosage forms can be a daily dose which is administered once a day or being divided to be administered several times a day, or alternative in a sustained release form. Suitable dosage forms are intended for oral administration.

All benzothiazepines, however, will not be effective as IBAT inhibitor compounds. Thus, diltiazem, which is a 1,5-benzothiazepine, is a calcium blocker with coronary vasodilating activity (see The Merck Index, Merck & Co, Inc., 12th ed., 1996, p. 541). With respect to inhibition of IBAT, diltiazem has no activity.

5

In general, pharmaceutical drug substances will be absorbed in the upper small intestine, and therefore only a small amount will reach ileum when administered in a conventional oral dosage form. Irrespective of the construction of the pharmaceutical dosage form, it should provide contact for the active compound, e.g. inhibitor of IBAT, with the compound's site of action in the body, for example in the ileum. The above prior art documents discuss in general terms suitable pharmaceutical dosage forms for the described IBAT inhibitor compounds. However, none of the documents describe a specific way to obtain a release of the active substance directly to or close to the site of action.

15 Contact between the active drug and the site of action can be established in different ways. The present application describes a new pharmaceutical dosage form which reduces and minimises absorption, metabolism and dilution in the luminal content of the IBAT inhibitor in the body before the active substance (IBAT inhibitor compound) reaches its site of action.

20

It has been proposed that after absorption over the gastro-intestinal membrane, an IBAT inhibitor could interact with transport systems similar to IBAT for instance the corresponding transport system in the liver (LBAT) or could provide other non-specific systemic effects which could lead to undesirable pharmacological or even toxicological effects. This could severely limit the clinical usefulness of IBAT inhibitors especially in the treatment of hypercholesterolaemia, i.e. conditions associated with elevated concentrations of total cholesterol and low-density lipoprotein cholesterol.

25 The inhibition of the re-absorption of bile acids from the small intestine performed by an effective IBAT inhibitor may lead to increased levels of bile acids in the lower parts

30

(colon) of the gastro-intestinal tract. Such an increase of bile acid concentrations in the distal regions could potentially generate diarrhoea and discomfort to the patient. The present invention provides a new approach to minimise the concentration of free bile acids in the colon and thereby reduce the potential risk of adverse events by co-administration of a bile acid binder together with the IBAT inhibitor. However, the combination of an IBAT inhibitor and a bile acid binder have previously been proposed in the above patent applications describing new IBAT inhibitor compounds. The purpose of such previously described combinations have been to enhance the cholesterol lowering efficacy of the therapy, and there is no hint that such a combination could be used to minimise a potential risk for diarrhoea connected with IBAT inhibitor therapy.

BRIEF DESCRIPTION OF THE INVENTION

The aim of the present invention is to reduce the problem with undesirable side effects of IBAT inhibitor compounds by providing a pharmaceutical formulation, which reduces the systemic drug exposure while maintaining or enhancing the cholesterol lowering effect of the drug. Such undesirable systemic effects put a load on other organs, e.g. liver and kidneys. Thus, the present dosage form provides a reduced, i.e. minimum absorption, metabolism and dilution in the luminal content of the IBAT inhibitor by a specific targeting to the site of action. The release is directed specifically to the site of action which reduces or even might avoid toxicological effects of the drug. The formulation is intended to be orally administered and pass through the upper part of the small intestine with a minimum release of the IBAT inhibitor before it reaches the distal jejunum or proximal ileum.

The present invention provides such a dosage form, which delivers the main part of the dose to the site of action, i.e. in the distal jejunum, in the proximal ileum or in the distal ileum. The release of the drug is thereby reduced or minimised to more proximal parts, the duodenum and jejunum, where drug absorption in general is most efficient. Thus, the

release of the drug should preferably start in the distal jejunum or proximal ileum, or the entire dose should be delivered directly to the ileum.

Preferably, the formulation is an orally administered formulation, such as a delayed release
5 formulation, which starts to release the main part of the drug in the distal jejunum or in the proximal ileum. The oral formulation might also provide protection of the drug from the acid environment in the stomach by an enteric coating. Such an enteric coating also protects the gastric mucosa from drug exposure and thereby minimises irritation or even damages of the gastric mucosa potentially caused by aggressive drug exposure.

10

An additional aim of the present invention is to provide a combination for simultaneous, separate or sequential administration which combination comprises an IBAT inhibitor and a bile acid binder. Such a combination will protect the patient from any possible side effect caused by excess of bile acids in the colon, such as diarrhoea. If the transport of bile acids
15 is blocked by an IBAT inhibitor the bile acids might be deposited in the colon and induce a secretory diarrhoea - by irritation and inflammation - as a undesired side effect caused by the treatment with an IBAT inhibitor.

Another aspect of the provided combination therapy is that the bile acid binder, for
20 instance a resin such as cholestyramine or cholestipol, could preferably be administered in a dosage form with colon release of the bile acid binder. A colon release formulation will provide protection of the bile acid binder to the luminal contents in the more proximal parts of the intestine, where the bile acid concentrations are high. Such a formulation will prevent binding of bile acids to the bile acid binder before the formulation reaches the
25 colon. Thereby, maximal bile acid binding capacity will be obtained in the colon and any possible gastro-intestinal side effects, such as diarrhoea, may be avoided. Thus, any additional amount of bile acid presented in the colon due to the treatment with the IBAT inhibitor compound, would be bound to a bile acid binder, which the bile acid binder is preferably delivered in the colon, thereby any possible side effects such as diarrhoea is
30 avoided.

DETAILED DESCRIPTION OF THE INVENTION

IBAT inhibitor compounds

5

Active ingredients suitable as IBAT inhibitor compounds in the present invention are those exhibiting activity when screening for IBAT inhibiting properties. Suitable examples of such compounds can be found in the references cited on page 2 of the present application.

- 10 Active ingredients particularly suitable as IBAT inhibitor compounds in the present invention include benzothiazepines, and more particularly 1,4-benzothiazepines and 1,5-benzothiazepines exhibiting activity when screening for IBAT inhibiting properties. Of these, compounds with an oxidized sulfur group, particularly a sulfone group, in the 7 membered ring are preferred. Furthermore, the presence of an amine group in the 7
- 15 membered ring is preferred.

Pharmaceutical formulations

- According to one aspect of the invention, an orally administered pharmaceutical formu-
- 20 lation of an IBAT inhibitor compound is provided, which formulation releases almost the entire dose of the IBAT inhibitor compound in the distal jejunum, in the proximal ileum, or deliver the dose directly to the ileum. Such a formulation will minimise drug release in the upper part of the small intestine, i.e. above distal jejunum.

- 25 Optimal drug release and drug binding in the ileum can for instance be obtained by a delayed release formulation, such as a formulation with a specified lagtime. More specifically, less than 30 % of the drug could be released during the time the formulation spends in the stomach and in the proximal small intestine, i.e. during the passage of the upper part of the small intestine.

30

Thus, according to a second aspect, the present invention provides a pharmaceutical formulation with a delayed release of the IBAT inhibitor compound by a controlled lagtime. The main part of the formulation shall pass the duodenum and jejunum with a minimum release of the active dose, and thereby increasing the dose available for binding to the site of action in the ileum and thereby increasing the inhibition of the ileal bile acid transport system. Preferably, the lagtime period is about 0.5 - 2 hours calculated from emptying from the stomach, and more than 70 % of the dose should be released approximately during the next 0.5 - 2.0 hours, i.e. after the lagtime period. More preferably, the dose should be released during the first hour after the lagtime period.

Dosage forms with a controlled lagtime can be constructed in different ways for instance as described in the following.

A controlled lagtime can be triggered by pH changes, redox potential differences or luminal metabolic changes in the gastro-intestinal tract as described in Aliment Pharmacol Ther 1997, 11 (suppl 3): 109-115. Such a controlled lagtime could be obtained for instance by a programmed disintegration of the formulation due to erosion, dissolution or in general by components present in the formulation interacting with the environment in the gastro-intestinal tract. Preferably, the drug release from the dosage form could be triggered by the pH variation between jejunum and ileum.

Alternatively, the drug release from the dosage form can be chronographic controlled to obtain the above specified time limits, such as for instance described in the European Patent Application, Publication No. EP-A-0384642.

When the formulation reaches the distal jejunum or the ileum, the drug release should preferably be either immediately, with a sustained release or be based on a combination of such release principles. The duration of the drug release for a sustained release formulation should preferably not exceed 2 hours.

According to a third aspect of the invention, a sustained release formulation can be constructed by any known principle, such as eroding or non-eroding matrices, membrane-coating layers or by diffusion or osmotically driven drug release. Suitable techniques for the construction of such formulations are for instance described in M. E. Aulton,
5 Pharmaceutics, The science of dosage form design. (1988).

An additional aspect of the invention is to combine an IBAT inhibitor compound with a bile acid binder thereby avoiding a possible risk of excess of bile acids in colon caused by the inhibition of the ileal bile acid transport system. An excess of bile acids in the visceral
10 contents may cause diarrhoea. Thus, the present invention also provides a treatment of a possible side effect such as diarrhoea in patients during therapy comprising IBAT inhibitor compounds.

Suitable bile acid binders for such a combination therapy are resins, such as cholestyrmine
15 and cholestipol. One advantage is that the dose of bile acid binder might be kept lower than the therapeutical dose for treatment of cholesterolemia in single treatment comprising solely a bile acid binder. By a low dose of bile acid binder any possible side effects caused by poor tolerance of the patient to the therapeutic dose could also be avoided.

A further aspect in connection with such a combination therapy is that the bile acid binder could be administered in a dosage form with colon release, i.e. delivery of the active dose of bile acid binder in the colon. A possible risk of receiving an excess of bile acid in the colon by treatment with an IBAT inhibitor could be avoided by co-administration of a bile acid binder with colon release. Thus, any excess of bile acid in the colon, with a possible
25 risk to cause diarrhoea, will be bound into a resin. The dose of the bile acid binder could be kept low due to an effective use of the dose by such a colon release. The colon delivery of the bile acid binder can be obtained by a formulation comprising a core containing the bile acid binder and optionally pharmaceutically acceptable excipients, and a coating of said core with a delayed release membrane adapted for colonic delivery. Technologies to obtain

such a delivery of drugs to the colon are for example described in Drug Development and Industrial Pharmacy 1997, 23: 893-913.

Further general aspects of the invention are that the formulations can be solid, semi-solid
5 or liquid formulations. In a solid formulation, the carrier can be monolithic, such as tablets or capsules. One preferred monolithic formulation is a coated tablet, a capsule comprising small, coated units or a multiple unit tablet comprising a multitude of small coated units. Semi-solid or liquid formulations can be administered in capsules suitable for such vehicles. The most preferred formulation is an oral formulation such as a tablet or a
10 capsule comprising coated small units or pellets. The formulation or dosage form may contain from 0.05% to 95% of the active compound in admixture with a pharmaceutically acceptable carrier, or pharmaceutically acceptable excipients.

Preparation of core material

15

The core material for the units, i.e. the tablets or the individual pellets can be constituted according to different principles. The core material may be homogenous or heterogeneous. The core containing the active principle may be differently formulated such as monolithic tablets, capsules, granules, pellets, other particles or crystals.

20

With a homogenous core material is meant, that it has a homogenous distribution of active substance throughout the core material.

The active substance, i.e. the IBAT inhibitor compound, is optionally mixed with further
25 components to obtain preferred handling and processing properties and a suitable concentration of the active substance in the final mixture. Such components can be binders, surfactants, lubricants, glidants, fillers, additives or other pharmaceutically acceptable ingredients, alone or in mixtures.

Said core material may be produced either by direct compression of the mixed ingredients, or by granulation of the ingredients followed by compression of the granulated material.

In direct compression, the ingredients are mixed and compressed by using ordinary tableting equipment.

5

For the granulation there are numerous alternatives of granulating procedures mentioned in the literature, dry methods like roller compaction (Chilsonator) and wet methods utilizing granulating solutions with and without the addition of binders. A variant of the wet methods is to make a spray-granulation in a fluid bed.

10

For the wet granulating methods, either organic solvents, aqueous solutions or pure water may be utilized to prepare the granulating solutions. Due to environmental considerations pure water is preferred, if it is possible due to the composition of the mixture.

15 Homogenous core particles can also be prepared by techniques such as dry or wet milling, freeze milling, air-jet micronisation, spray drying, spray chilling, controlled crystallisation, supercritical crystallisation, emulsion solvent evaporation and emulsion solvent extraction.

The core material may also be produced by extrusion/spheronization, balling or
20 compression, utilizing different process equipments.

The size of the formulated core materials is approximately between 2 and 14 mm, preferably between 3 and 9 mm for a tablet preparation, and between 0.001 and 4 mm, preferably between 0.001 and 2 mm for a pellet preparation.

25

The manufactured core material may be further layered with additional ingredients comprising the active substance and/or be used for further processing.

Alternatively, the core material may be heterogeneous with an inner zone, for instance a
30 seed or sphere, not containing the active substance. A layer comprising the active

substance, and optionally pharmaceutically acceptable excipients, surrounds this seed or sphere.

The seed or sphere may be soluble or insoluble. Optionally, the seed or sphere (inner zone)
5 may be coated with an inert layer to prepare a smooth surface before the layer containing active substance is applied onto the seed/sphere.

Insoluble seeds/spheres may comprise different oxides, celluloses, organic polymers and other materials, alone or in mixtures. Water-soluble seeds/spheres may comprise different
10 inorganic salts, sugars and other materials, alone or in mixtures. The size of the seeds may vary between approximately 0.1 and 2 mm. The seeds layered with the matrix containing the active substance are produced either by powder or solution/suspension layering using for instance granulating or spray coating/layering equipment.

15 *Processes for application of delayed release membranes*

Delayed release membrane can be applied to the core material, being a monolithic tablet, multiple units or a hard or soft gelatine capsule, by coating or layering procedures in suitable equipment such as coating pans, coating granulators or in a fluidized bed apparatus
20 using water and/or organic solvents for the coating process. Also powder-coating principles may be applied. Another possibility is to apply the coating by microencapsulation techniques such as coacervation, emulsification with subsequent removal of the solvent by extraction or evaporation, ionotropic gelation or congealing.

25 Such delayed release membranes may be applied on core material comprising the IBAT inhibitor for delivery to the distal small intestine and optionally also be applied to the bile acid binder for delivery to the colon.

Pharmaceutical additives

Delayed release coatings may be obtained by one or more, separately or in compatible combinations of pharmaceutically acceptable ingredients, in amounts carefully titrated to reach the intended release properties. As coating layer, the following pH sensitive polymers can be applied; e.g. methacrylic acid copolymers, cellulose acetate phthalate, hydroxypropyl methylcellulose phthalate, hydroxypropyl methylcellulose acetate succinate, polyvinyl acetate phthalate, cellulose acetate trimellitate, carboxymethyl ethyl-cellulose, shellac or other suitable enteric coating layer polymer(s). The coating layer may also be composed of film-forming polymers being sensitive to other luminal components than pH, such as bacterial degradation or a component that has such a sensitivity when it is mixed with another film-forming polymer. Examples of such components providing delayed release to the intended regions are; polymers comprising azo bond(s), polysaccharides such as pectin and its salts, galactomannans, amylose and chondroitin, disulphide polymers and glycosides.

The delayed release coating or an additional coating of the formulation may contain other film-forming polymers being non-sensitive to the luminal conditions for technical reasons or chronographic control of the drug release. Materials to be used for such purpose includes, but are not limited to; sugar, polyethylene glycol, polyvinylpyrrolidone, polyvinyl alcohol, polyvinyl acetate, hydroxypropyl cellulose, methylcellulose, ethylcellulose, hydroxypropyl methylcellulose, carboxymethylcellulose sodium and others, used alone or in mixtures.

Additives such as dispersants, colorants, pigments, additional polymers e.g. poly(ethylacrylat, methylmethacrylat), anti-tacking and anti-foaming agents may also be included into the coating layer. Other compounds may be added to increase film thickness and to decrease diffusion of acidic gastric juices into the core material.

The coating layers may also contain pharmaceutically acceptable plasticizers to obtain desired mechanical properties. Such plasticizers are for instance, but not restricted to, triacetin, citric acid esters, phthalic acid esters, dibutyl sebacate, cetyl alcohol, polyethylene glycols, glycerol monoesters, polysorbates or other plasticizers and mixtures thereof.

- 5 The amount of plasticizer is preferably optimised for each formula, in relation to the selected polymer(s), selected plasticizer(s) and the applied amount of said polymer(s).

- In preparation of tablets, either as monolithic drug containing cores for subsequent coating with a delayed release membrane or as a matrix for coated multiple units, additional
- 10 ingredients may be needed to obtain suitable technical properties such as binders, disintegrants, bulk agents, glidants, lubricants, and coatings agents without effects on the drug release such as water soluble polymers, anti-tacking agents, colourants, pigments and waxes. Ingredients well known for such usage are for example described in "Handbook of pharmaceutical excipients", 2nd edition, 1994, Pharmaceutical Press, London.

15

Preparation of final dosage forms

- Coated units may be filled into hard gelatine capsules or mixed with tablet excipients, such as fillers, binders, disintegrants, lubricants and other pharmaceutically acceptable addi-
- 20 tives, and be compressed into tablets. The compressed tablet is optionally covered with film-forming agents to obtain a smooth surface of the tablet and further enhance the mechanical stability of the tablet during packaging and transport. Such a tablet coat, which may be applied on a multiple unit tablet or a conventional tablet, may further comprise additives like anti-tacking agents, colourants and pigments or other additives to improve
- 25 the tablet appearance.

Suitable drugs for the new formulations are IBAT inhibitor compounds such as described in the above-discussed documents, hereby incorporated by references.

The IBAT inhibitor compound could alternatively be a low permeability drug as defined in the Biopharmaceutical Classification System proposed by FDA.

5 A combination therapy according to the invention should preferably comprise simultaneously, separately or sequentially administration of an IBAT inhibitor compound and a bile acid binder. The IBAT inhibitor could preferably be formulated for ileum delivery and the bile acid binder could preferably be formulation for colon release.

Medical and pharmaceutical use of the invention

10

The pharmaceutical formulations according to the present invention can be used in the treatment of hypercholesterolaemia. A suitable unit dose will vary with respect to the patients body weight, condition and disease severity. The dose will also depend on if it is to be used for prophylaxis or in the treatment of severe conditions, as well as the route of
15 administration. The daily dose can be administered as a single dose or divided into two or more unit doses. An orally administered daily dose of an IBAT inhibitor is preferably within 0.1 – 1,000 mg, more preferable 1 - 100 mg.

20 A pharmaceutical formulation according to the present invention with a targeted delivery in the gastro intestinal tract provides a reduced systemic exposure, as can be measured by the area under the drug plasma concentration versus time curve (AUC), while maintaining or even increasing the therapeutic effect, as e.g. measured by serum cholesterol reduction.

25 A combination therapy comprising an IBAT inhibitor and a bile acid binder comprises preferably a low daily dose of the bile acid binder, such as less than 5 g of a resin, and more preferably less than 2 g. A dosage form with colon release of the bile acid binder could be constructed by any of the above described principles for delayed release formulations.

The following contemplated Examples are intended to illustrate, but in no way limit the scope of the invention.

EXAMPLES

5 Example 1

A formulation having the following composition can be prepared:

	amount/capsule (mg)
10 IBAT inhibitor compound	
(1,5-benzothiazepine)	10
Non pareil spheres	500
Ethyl cellulose	2
Hydroxypropylmethyl cellulose	10
15 Eudragit L100-55	25
Triethylcitrate	2.4

The active drug can be dissolved together with ethyl cellulose and hydroxypropyl cellulose in ethanol 99 %. The mixture can then be sprayed onto the non-pareil spheres in a fluidized
20 bed apparatus. Thereafter, the pellets can be dried and aerated to remove residual ethanol. The Eudragit L100-55 dispersion with addition of triethyl citrate can then be sprayed onto the drug beads in a fluidized bed apparatus. Subsequently, the coated beads can be filled in hard gelatine capsules after drying and sieving.

25 Example 2

A formulation having the following composition can be prepared:

	amount/tablet (mg)
IBAT inhibitor compound	
(1,5-benzothiazepine)	10
Silicon dioxide	200
5 Povidone K-25	20
Eudragit FS30D	30
Microcrystalline cellulose	250
Sodium stearyl fumarate	5

10 The active drug can be suspended in water and sprayed onto silicon dioxide cores of a predefined size in a fluidized bed apparatus. The drug pellets can be dried in an oven at 40° C for 24 h. Thereafter, a layer of Povidone K-25 can be applied on the beads from an ethanolic solution in a fluidized bed apparatus. A final coat of Eudragit FS30D dispersion can be applied thereafter in a fluidized bed. The coated beads can be mixed with

15 microcrystalline cellulose and sodium stearyl fumarate in a mixer and subsequently compressed to tablets.

CLAIMS

1. An oral pharmaceutical formulation comprising an inhibitor compound of the ileal bile
5 acid transport (IBAT inhibitor compound) and a pharmaceutically acceptable carrier,
wherein the formulation is designed to deliver the IBAT inhibitor compound in the
ileum.
2. The oral pharmaceutical formulation according to claim 1, wherein the formulation is
10 designed to deliver the IBAT inhibitor compound in the ileum by release in one or
more parts of the body selected from the distal jejunum and proximal ileum, and/or
directly in the ileum.
3. The formulation according to claim 1, wherein the carrier is designed to deliver the
15 IBAT inhibitor compound in the ileum.
4. The formulation according to claim 1, wherein the carrier is designed to release the
IBAT inhibitor compound in the distal jejunum and in the proximal ileum.
- 20 5. The formulation according to any one of the claims 1 to 4, wherein the carrier is
designed to give a minimum release of the IBAT inhibitor compound in the upper part
of the small intestine.
6. The formulation according to any one of claims 1 to 4, wherein the pharmaceutical
25 formulation is a delayed release formulation.
7. The formulation according to claim 6, wherein the formulation provides a lagtime of
about 0.5 - 2 hours after emptying the stomach.

8. The formulation according to claim 7, wherein the IBAT inhibitor compound is released during the first hour after the lagtime.
9. The formulation according to claim 6, wherein release of the IBAT inhibitor compound from the delayed release formulation is triggered by the pH differences between the jejunum and ileum.
10. The formulation according to any one of claims 1 to 9, wherein the IBAT inhibitor compound is a low permeability drug as defined in the Biopharmaceutical Classification System FDA.
11. The use of a pharmaceutical formulation comprising an IBAT inhibitor compound with targeted delivery in the gastro-intestinal tract according to any one of the claims 1 to 10 to reduce systemic exposure.
12. The use of a pharmaceutical formulation comprising an IBAT inhibitor compound with targeted delivery in the gastro-intestinal tract according to any one of the claims 1 to 10 to enhance the therapeutic effect.
13. The use of a pharmaceutical formulation according to any one of the claims 1 to 10 in the treatment of hypercholesterolemia.
14. The use of a pharmaceutical formulation according to any one of the claims 1 to 10, in the manufacture of a medicament for the prophylactic or therapeutic treatment of hypercholesterolemia.
15. A method for prophylactic or therapeutic treatment of a subject suffering from, or susceptible to, hypercholesterolemia, which method comprises administering to the subject a pharmaceutical formulation designed according to any one of claims 1 to 10.

16. A pharmaceutical formulation for simultaneous, separate or sequential administration in the prophylactic or therapeutic treatment of hypercholesterolemia, which formulation comprises an IBAT inhibitor compound and a bile acid binder.
- 5 17. The pharmaceutical formulation according to claim 16, wherein the IBAT inhibitor compound is a low permeability drug as defined in claim 10.
18. The pharmaceutical formulation according to claim 16, wherein the bile acid binder is a resin.
- 10 19. The pharmaceutical formulation according to claim 18, wherein the bile acid binder is in a formulation with colon release.
20. The use of a pharmaceutical formulation according to any one of claims 16 - 19 in the treatment of diarrhoea during therapy comprising an IBAT inhibitor compound.
- 15 21. The use of a pharmaceutical formulation according to any one of the claims 16 to 20, in the manufacture of a medicament for the prophylactic or therapeutic treatment of hypercholesterolemia.
- 20 22. A method for prophylactic or therapeutic treatment of a subject suffering from, or susceptible to, diarrhoea during therapy comprising an IBAT inhibitor compound, which method comprises administering to the subject a pharmaceutical formulation designed according to any one of claims 15 to 18.
- 25 23. The use of a bile acid binder as prophylaxis or in the treatment of diarrhoea during therapy comprising an IBAT inhibitor compound.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 00/00755

A. CLASSIFICATION OF SUBJECT MATTER

IPC7: A61K 45/00, A61K 9/20, A61P 3/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: A61K, A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5614220 A (YOSHIYUKI HIRAKAWA ET AL), 25 March 1997 (25.03.97) --	1-23
X	US 5723458 A (LAWRENCE EDWARD BRIEADDY ET AL), 3 March 1998 (03.03.98) --	1-23
A	Dialog Information Services, File 73, EMBASE, Dialog accession no. 02888148, Embase accession no. 1985132107, Jacobsen O. et al: "Effect of entercoated cholestyramine on bowel habit after ileal resection: A double blind crossover study"; & British Medical Journal (BR. MED. J.) (United Kingdoms) 1985, 290/6478 (1315-1318) -- -----	1-23

☐ Further documents are listed in the continuation of Box C.
 ☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

31 July 2000

Date of mailing of the international search report

03-08-2000

 Name and mailing address of the ISA/
 Swedish Patent Office
 Box 5055, S-102 42 STOCKHOLM
 Facsimile No. +46 8 666 02 86

Authorized officer

 Anneli Jönsson/gh
 Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE00/00755

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 11-15, 20, 22-23
because they relate to subject matter not required to be searched by this Authority, namely:
see next sheet
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).:

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE00/00755

Claims 11-15,20,22-23 relate to methods of treatment of the human or animal body by surgery or by therapy/diagnostic methods practised on the human or animal body/ Rule. 39.1.(iv). Nevertheless, a search has been executed for these claims. The search has been based on the alleged effects of the compounds/compositions.

INTERNATIONAL SEARCH REPORT
Information on patent family members

02/12/99

International application No.

PCT/SE 00/00755

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5614220 A	25/03/97	CA 2142142 A	11/08/95
		CN 1116926 A	21/02/96
		EP 0671167 A	13/09/95
		JP 7223970 A	22/08/95

US 5723458 A	03/03/98	AU 6008894 A	29/08/94
		CA 2156183 A	18/08/94
		EP 0683774 A	29/11/95
		HU 71610 A	29/01/96
		HU 9501818 D	00/00/00
		IL 108633 A	15/07/98
		JP 2886341 B	26/04/99
		JP 8506576 T	16/07/96
		WO 9418184 A	18/08/94
		ZA 9401003 A	14/08/95
GB 9400314 D	09/03/94		
